

**Desmond, E.** (2006): Reducing salt: A challenge for the meat industry. *Meat Science*, 74, 188-196.

**FSA.** (2006): Salt reduction targets. Available from: <http://www.food.gov.uk/multimedia/pdfs/salttargetsapril06.pdf>.

**FSAI.** (2005): Salt and health: review of the scientific evidence and recommendations for public policy in Ireland. Food Safety Authority of Ireland.

**Gelabert, J., P. Gou, L. Guerrero, J. Arnau** (2003): Effect of sodium chloride replacement on some characteristics of fermented sausages. *Meat science*, 65, 833-839.

**Gimeno, O., I. Astiasarán, J. Bello** (1998): A mixture of potassium, magnesium and calcium chlorides as a partial replacement of sodium chloride in dry fermented sausages. *Journal of Agricultural and Food Chemistry*, 46, 4372-4375.

**Gimeno, O., I. Astiasarán, J. Bello** (1999): Influence of partial replacement of NaCl with KCl and CaCl<sub>2</sub> on texture and color of dry fermented sausages. *Journal of Agricultural and Food Chemistry*, 47, 873-877.

**Gimeno, O., I. Astiasarán, J. Bello** (2001): Calcium ascorbate as a potential substitute for NaCl in dry fermented sausages: effect on colour, texture and hygienic quality at different concentrations. *Meat Science*, 57, 23-29.

**Gou, P., L. Guerrero, J. Gelabert, J. Arnau** (1996): Potassium chloride, potassium lactate and glycine as sodium chloride substitutes in fermented sausages and in dry-cured pork

loin. *Meat Science*, 42, 37-48.

**Guàrdia, M. D., L. Guerrero, J. Gelabert, P. Gou, J. Arnau** (2006): Consumer attitude towards sodium reduction in meat products and acceptability of fermented sausages with reduced sodium content. *Meat Science*, 73, 484-490.

**Guàrdia, M. D., L. Guerrero, J. Gelabert, P. Gou, J. Arnau** (2008): Sensory characterisation and consumer acceptability of small calibre fermented sausages with 50% substitution of NaCl by mixtures of KCl and potassium lactate. *Meat Science*, 80, 1225-1230.

**Ibáñez, C., L. Quintanilla, A. Irigoyen, I. García-Jalón, C. Cid, I. Astiasarán, J. Bello** (1995): Partial replacement of sodium chloride with potassium chloride in dry fermented sausages: influence on carbohydrate fermentation and nitrosation process. *Meat Science*, 40 (1), 45-53.

**ISO Method 1841** (1970): Determination of salt. International Standards Meat & Meat products. Ginebra. International Organization for Standardization.

**ISO Method 2917** (1999): Measurement of pH. International standards meat and meat products. Genève Switzerland: International Organisation for Standardisation.

**Muguerza, E., O. Gimeno, D. Ansorena, I. Astiasarán** (2004): New formulations for healthier dry fermented sausages: a review. *Trends in Food Science & Technology*, 15, 452-457.

**Ockerman, H. W., L. Basu** (2007): Produc-

tion and consumption of fermented meat products. In F. Toldá (Ed.), *Handbook of fermented meat and poultry* (pp. 9-15). Iowa, USA: Blackwell Publishing.

**Pravilnik o mikrobiološkim kriterijima za hranu** (NN 74/08, NN 156/08, NN 89/10)

**Ruusunen M., E. Puolanne** (2005): Reducing sodium intake from meat products. *Meat Science*, 70, 531-541.

**Stahkne, L. H., K. Tjener** (2007): Influence of processing parameters on cultures performance. In F. Toldá (Ed.), *Handbook of fermented meat and poultry* (pp. 187-194). Iowa, USA: Blackwell Publishing.

**Vodič za mikrobiološke kriterije za hranu.** Ministarstvo poljoprivrede, ribarstva i ruralnog razvoja. Treće izmjenjeno izdanje. Zagreb, 2011.

**Toldrá, F.** (2007): *Handbook of Fermented Meat and Poultry*. Oxford OX4 2DQ, UK: Blackwell Publishing Ltd.

**US Department of Health and Human Services** (2005). 2005 Dietary guidelines for Americans. Available from: <http://www.health.gov/dietaryguidelines/dga2005/document>.

**Zanardi, E., S. Ghidini, M. Conter, A. Ianieri** (2010): Mineral composition of Italian salami and effect of NaCl partial replacement on compositional, physico-chemical and sensory parameters. *Meat science*, 86, 742-747.

Received: June 1, 2011

Accepted: July 5, 2011

**Primošten/Croatia, 19-22 October 2011**

**Power of Fungi and Mycotoxins in Health and Disease**

**General Information**

- Venue & Accommodation:** Hotel Zora, Radula 11, HR-22202 Primošten, Croatia. T: +385 22 581 009, F: +385 22 571 120, [www.zora-hotel.com](http://www.zora-hotel.com)
- Date:** Wednesday, October 19 till Saturday, October 22, 2011
- Symposium Language:** English
- Meeting Secretariat:** Schrotova 39, HR-10000 Zagreb, Croatia. T: +385 1 6394 493, F: +385 1 6394 494, [info@hmd-cms.hr](mailto:info@hmd-cms.hr), [www.hmd-cms.hr](http://www.hmd-cms.hr)
- Organised by:** Croatian Microbiological Society
- Supported by:** Federation of European Microbiological Societies (FEMS)

## The quality of meat during t

### Summary

*Production of fermented sausages of horse meat is a traditional manufacture in some regions. The aim of this study was to investigate the physicochemical and sensory properties of sausages made from horse meat, depending on the season of production and water content, water activity 0.896, the average values of pH in the finished product amounted to 5.83, 19.90% NaCl 3.44% and ammonia 0.42 mg%. Slight differences were detected in sensory properties of sausages in relation to the production season (p < 0.05) in physicochemical properties. Key words: fermented sausages, horse meat.*

### Introduction

As a commercially most valuable sausage manufacturing product, fermented sausages are manufactured of the finest lots of meat of various animal species. In that sense, horse meat, according to its chemical composition and manufacturing values, is a very convenient substrate for the production and obtained fermented sausage is a product of extraordinary quality and exceptional value (Feiner, 2006; Dobranić et al., 2008, 2009). Production of horse meat and horse meat products is specific to individual countries and regions, while in some parts of the world and cultures it is not present at all. Dur-

- <sup>1</sup> Damir Alagić, PhD, College of Agriculture at Križevci
- <sup>2</sup> Nevijo Zdolec, PhD, scientific novice - assistant, Fleck, PhD, assistant professor; Lidija Kozačinskić, PhD, assistant professor; Lidija Kozačinskić, PhD, assistant professor; Lidija Kozačinskić, PhD, assistant professor
- <sup>3</sup> Anamaria Ekert Kabalin, PhD, assistant professor
- <sup>4</sup> Gorjarić Alagić, dipl.ing.preh.teh., Trade
- <sup>5</sup> Miomir Stojnović, MSc, College of Agriculture at

## mented sausages of horse meat during the three production seasons

Alagić, D.<sup>1</sup>, N. Zdolec<sup>2</sup>, B. Njari<sup>2</sup>, I. Filipović<sup>2</sup>, A. Ekert-Kabalin<sup>3</sup>, G. Čorić-Alagić<sup>4</sup>, M. Stojnović<sup>5</sup>, Ž. Cvrtlić-Fleck<sup>2</sup>, L. Kozačinskić<sup>2</sup>

scientific paper

*this country has a long tradition that enriches our cuisine. However, there is little and product quality to be able to talk about standardization and protection of the physicochemical and sensory properties of sausages made from horse meat, depending on the season of production and water content, water activity 0.896, the average values of pH in the finished product amounted to 5.83, 19.90% NaCl 3.44% and ammonia 0.42 mg%. Slight differences were detected in sensory properties of sausages in relation to the production season (p < 0.05) in physicochemical properties. Key words: fermented sausages, horse meat.*

2005, eight major horse meat processing countries produced over 100 tons (FAO-UN Horticultural base). In Croatia, this activity is related to several regional manufacturers, which is probably not intended to exploit the potential of horse meat as a quality raw material for the development of specific products in the indigenous cuisine (Njari, 2010). In the villages of Pakrac, a tradition of sausage production from horsemeat was brought from their original homeland and revived by the Italian minority. In produce a homemade durable product, horse meat sausage (ta"). In this area in the past few

years there were mainly slaughtered about 1500 horses. Also, in the area of Varazdin there is a long tradition of production of sausage products made of horse meat (registered sausage- butcher's trade Zvonko Šanjek Family Crafts since 1954). Simić and Mioković (2008) point out that horse sausages in our country are produced in homes or small crafts, so there is little data on their composition and quality. Thus, Dević and Stamenković (1990) explored the possibility of producing horsemeat sausages cooked and smoked with hot smoke, then dried. They believe that these are the sausages of acceptable organoleptic properties and that

merca 1, 48260 Križevci

<sup>1</sup> PhD, full professor, Ivana Filipović, univ.mag.spec., DVM, scientific novice - assistant, Željka Cvrtlić professor; University of Zagreb, Faculty of Veterinary Medicine, Dpt. of Hygiene and Technology of Foodstuffs

University of Zagreb, Faculty of Veterinary Medicine, Dpt. of Animal Husbandry, Heinzelova 55, Zagreb  
primnica, Trg Slobode 7, 48 000 Koprivnica  
1.Demerca 1, 48260 Križevci




Figure 1 Horse meat intended for fermented sausages production



Figure 2 Pork fat intended for fermented sausages production

Table 1 pH during maturing of horsemeat fermented sausages

Parameter	Batch	Days			
		0.	14.	28.	36.
		X ± SD	X ± SD	X ± SD	X ± SD
pH	April	5,82 <sup>a</sup> ± 0,01	5,36 <sup>a</sup> ± 0,02	5,04 <sup>a</sup> ± 0,01	5,87 <sup>a</sup> ± 0,01
	September	6,19 <sup>a</sup> ± 0,02	5,60 <sup>a</sup> ± 0,03	5,24 <sup>a</sup> ± 0,02	5,82 <sup>a</sup> ± 0,01
	November	6,09 <sup>a</sup> ± 0,01	5,02 <sup>bc</sup> ± 0,01	5,53 ± 0,02	5,81 <sup>c</sup> ± 0,03

<sup>abc</sup> within the same row values marked with the same letter are statistically significantly different (p<0.01)

Table 2 Water content during maturing of horsemeat fermented sausages

Parameter	Batch	Days			
		0.	14.	28.	36.
		X±SD	X±SD	X±SD	X±SD
Moisture (%)	April	62,08 <sup>a</sup> ±1,04	34,44 <sup>a</sup> ±0,93	25,31 <sup>a</sup> ±0,47	19,69 <sup>a</sup> ±0,74
	September	60,80 <sup>a</sup> ±0,69	36,11 <sup>a</sup> ±0,66	26,28 <sup>a</sup> ±0,29	20,17 <sup>a</sup> ±0,64
	November	62,29 <sup>a</sup> ±0,74	34,17 <sup>a</sup> ±0,60	25,66 <sup>a</sup> ±0,67	19,86 <sup>a</sup> ±1,37

<sup>abc</sup> within the same row values marked with the same letter are statistically significantly different (p<0.01)

Table 3 Water activity during maturing of horsemeat fermented sausages

Parameter	Batch	Days			
		0.	14.	28.	36.
		X±SD	X±SD	X±SD	X±SD
Water activity	November	0,936 <sup>a</sup> ±0,002	0,925 <sup>a</sup> ±0,002	0,899 <sup>a</sup> ±0,002	0,896±0,001

<sup>abc</sup> within the same row values marked with the same letter are statistically significantly different (p<0.01)

can lead to interest of a wider circle of consumers. Simić and Mioković (2008) reported that horse fat is not suitable for the production of horse meat sausage because of its bad smell and taste and it is perishable.

Recent microbiological study of fermented sausages of horse meat in Croatia has shown that there is room for the improvement and standardization of these products (Alagić et al., 2008; Markov et al., 2010; Alagić

et al., 2011, submitted). In this paper we present the results of research on physicochemical and sensory properties of horse meat sausage, depending on the season of production and stage of maturity.

## Material and methods

### Sausage production and sampling

Horse meat sausages were produced in Varazdin Crafts house by the standard procedure that it applies. Sausages were made from horse meat (75%, Figure 1), solid pork fat (25%, Figure 2), and added ingredients (salt, pepper, paprika, garlic, nitrite salt). After filling of the intestine and draining on a stick, followed by cold smoking for 2-3 days, the sausages were left to mature in fermentation chamber until the 36th day.

Three sausages were sampled on day 0, 14, 28 and 36 (Figure 3-5). At the beginning of the production process, raw materials were also sampled, i.e. horse meat and pork fat tissue. Samples were sent to the laboratory in a portable refrigerator (+ 4 °C). All samples were subjected to physical and chemical tests and sensory evaluation in triplicate.

## Physico-chemical and sensory tests


pH values were determined in meat extract by digital pH meter (WTW, Germany), and the amount of salt and water during the ripening of sausages according to standard AOAC (2002) methodology. For the determination of water activity (aw), the sample was previously fragmented, and according to the manufacturer's instructions for portable aw - meter (Rotronic, Switzerland), aw value was measured: the vessel for the sample was filled, placed in the cradle and the measuring probe was put on it. The amount of ammonia was determined by microdiffusion quantitative method by Schmidt (Zivkovic, 1986). The sample mass of 1 g is chopped up and concentrically distributed in the outer chamber of Conway diffusion dish and poured with potassium carbonate. The solution of boric acid mixed with the indicator is poured in the interior chamber. Ammonia released from the meat reacts with boric acid and converts to ammonium-borate, and the amount of released NH<sub>3</sub> is proved by titration with hydrochloric acid. The end of titration is reflected in the emergence of a stable reddish liquid titrated. The amount of ammonia in mg% was calculated using the formula: consumption of HCl x 0.17 x 100 = mg% NH<sub>3</sub>.

Sensory properties were evaluated according to proceedings of Cocolina et al. (2005).

## Statistical data analysis

Data were analyzed using statistical software Statistica 8 (StatSoft, 2008). After the primary statistical analysis normality of distribution was checked by Kolmogorov-Smirnov normality distribution test. To determine the significance of the difference values determined during the ripening of sausages (and after each season) one-way analysis of variance was used (One way ANO-

<b>1 Before the tasting:</b>		
a) Color grade		
poor	(1 to 10)	well
b) assessing the appearance of cross-section		
poor	(1 to 10)	well
c) assessing connection of stuffing (association of fat and muscle tissue)		
poor	(1 to 10)	well
<b>2 Rating fragrance of products</b>		
Is there an odor?		
	YES	NO
If YES:		
Describe		
Quit Evaluation		
If NO, to evaluate		
poor	(1 to 10)	well
<b>3 Taste the product and assess:</b>		
a) rancidity		
poor	(1 to 10)	well
b) quality of fat		
poor	(1 to 10)	well
c) acidity		
poor	(1 to 10)	well
d) juiciness		
poor	(1 to 10)	well
e) tenderness		
poor	(1 to 10)	well
f) general taste		
poor	(1 to 10)	well
<b>4 Assessment 10 minutes after tasting:</b>		
poor	(1 to 10)	well
<b>5 Evaluating the overall sensory impression:</b>		
poor	(1 to 10)	well



<sup>ab</sup> values marked with the same letter (at same day of maturing) are statistically significantly different (p<0.05)

Graph 1 pH during maturing of horsemeat fermented sausages through three seasons


VA) with Tukey-HSD test for starting post-hoc analysis.

## Results and discussion

The results are shown in tables 1-5 and figures 1-5. Tables (modified physicochemical parameters during ripening) have shown significant difference between the average values determined for individual days (but within the same series that month). The significance of differences between the individual series (months), keeping in mind the days, is indicated on the graphs.


The average pH value of horse meat and pork fat tissue amounted to 5.72 or 6.29. During ripening, the pH value of sausage filling was decreasing for the first and second series (April and September) to day 28, and in the third series (November) to day 14 (Table 1). The final pH of sausages, compared to baseline values, decreased slightly (about 0.2 units). However, by day 28 of sausage ripening there was a significant acidification, which is characteristic for horse meat sausage. Specifically, Feiner (2006) points out that horse-meat acidifies much more than beef or pork because of their naturally higher amounts of glycogen. In processing beef and pork, the pH of the fermented sausages decreases by about 0.15 to 0.3 pH units as a result of degradation of natural sugar (glycogen).

Applying only lean horse meat, stuffing acidification is much more evident and the pH decreased to 0.7 to 0.8 pH units, which in our case is confirmed, but until day 28 of ripening. The average pH in the finished product was 5.83, which is within the results of Palearia et al. (2003). On the other hand, Markov et al. (2010) in domestic fermentations. Fermented sausages of horsemeat determine pH of 4.94, which is significantly lower than in our study. The differences are probably caused by the duration of



a,b values marked with the same letter (at same day of maturing) are statistically significantly different ( $p < 0.05$ )

Graph 2 Water content during maturing of horsemeat fermented sausages through three seasons



a,b values marked with the same letter (at same day of maturing) are statistically significantly different ( $p < 0.05$ )

Graph 3 NaCl content during maturing of horsemeat fermented sausages through three seasons

ripening and other influential factors. Graph 1 shows that pH differed significantly by the individual days of ripening in relation to the production season. It is commonly found in traditional manufacturing because it is difficult to ensure a standard quality of raw materials and continuous microclimate conditions. Feiner (2006) notes that the main difficulty during the production of fermented sausages is a hardly predictable degree of acidification of lean meat.

The reason is that neither the initial pH value, nor the share of sugar in the meat are never equal, and therefore buffering system of amino acids is not present in the meat (even the same kind of meat).

The amount of water is properly and continuously decreased during ripening, regardless of the season, from the initial 60% to 20% in the finished product (Table 2, Figure 2). Alagić et al. (2008) warned that a too



Figure 3 Horsemeat fermented sausages after stuffing (day 0)



Figure 4 Horsemeat fermented sausages at the end of maturing

Table 4 NaCl content during maturing of horsemeat fermented sausages

Parameter	Batch	Days			
		0.	14.	28.	36.
		X±SD	X±SD	X±SD	X±SD
NaCl (%)	April	2,74 <sup>a</sup> ±0,07	3,00 <sup>a</sup> ±0,08	3,30 <sup>a</sup> ±0,06	3,46 <sup>a</sup> ±0,04
	September	2,71 <sup>a</sup> ±0,08	3,03 <sup>a</sup> ±0,03	3,18 <sup>a</sup> ±0,08	3,34 <sup>a</sup> ±0,05
	November	2,78 <sup>a</sup> ±0,06	3,01 <sup>a</sup> ±0,03	3,16 <sup>a</sup> ±0,08	3,52 <sup>a</sup> ±0,06

a,b,c within the same row values marked with the same letter are statistically significantly different ( $p < 0.01$ )

Table 5 Ammonia content during maturing of horsemeat fermented sausages

Parameter	Batch	Days			
		0.	14.	28.	36.
		X±SD	X±SD	X±SD	X±SD
Ammonia (mg%)	April	0,82 <sup>a</sup> ±0,04	1,57 <sup>a</sup> ±0,05	0,52 <sup>a</sup> ±0,01	0,32 <sup>a</sup> ±0,03
	September	0,56 <sup>a</sup> ±0,04	1,52 <sup>ab</sup> ±0,07	0,52 <sup>a</sup> ±0,02	0,62 <sup>ab</sup> ±0,01
	November	0,63 <sup>a</sup> ±0,05	1,45 <sup>bc</sup> ±0,05	0,52 <sup>b</sup> ±0,08	0,31 <sup>bc</sup> ±0,06

a,b,c within the same row values marked with the same letter are statistically significantly different ( $p < 0.01$ )

long duration of horse sausage drying (42 days) results in a too dry and too hard final product (toughness, hard consistency, water content <20%). Markov et al. (2010) found twice as high water content in the domestic horse meat sausage. Studies of the chemical composition of fermented sausage meat from other species showed an average value of 25% (Zdolec et al. 2007; Pleadin et al., 2009).

Water activity was measured only in the third series of sausages, where we recorded an average value of

0.896 (Table 3), and Paleari et al. (2003) state that fermented products of horse meat have about 0.94 water activity. By reducing the amount of water, the amount of salt is increased from the initial 2.7 to 3.3 to 3.5% in all production batches (Table 4). The content of salt is consistent with the results reported by other authors (Kožačinski et al., 2008; Markov et al., 2010). The amount of ammonia in all production batches doubled until day 14 of ripening, after which there was a decrease towards the end of its maturity (Table 5). Sausages produced in September to the end of

ripening had a significantly higher amount of ammonia in comparison with the sausages from the season in April and November ( $p < 0.05$ ; graph 4). The research on fermented sausages showed that intense proteolytic processes occurred in the initial stages of ripening, which can be explained by activation of tissue proteases due to a lower pH (Hughes et al., 2002; Zdolec et al., 2008). Among other compounds, ammonia is a product of proteolysis, so our results were consistent with the foregoing. Sensory evaluation detected slight differences in sensory properties of sausages in relation to the season of production (Figure 5). The properties with the highest scores were connection of filling, the quality of fat and acidity, and the lowest properties of color, layout section, tenderness and juiciness.

In conclusion, based on the results of sensory index, we can conclude that the physicochemical changes in horse sausage filling during ripening resulted in quality products. Looking at seasonally-dependent movement of certain parameters, there are significant differences during ripening, which, however, did not significantly affect the uniformity of quality of finished products.


\* The results are a part of the PhD thesis of Damir Alagić "Factors of



fermentation in sausages filling from horse meat" Mentor: prof. PhD. Lidija Kozačinski (dissertation was prepared under the project of the Ministry of Science, Education and Sports of the Republic of Croatia 053-0531854-1853). We thank Mr. Zvonko Šanjek, the owner of sausage butcher's trade from Varazdin, who enabled us with his contribution to realize this work.


## References

- Alagić, D., L. Kozačinski, I. Filipović, N. Zdolec, M. Hadžiosmanović, B. Njari, Z. Kozačinski, S. Uhić (2008): Mikrobiološke promjene tijekom zrenja fermentiranih kobasica tijekom zrenja. Meso 10, 200-203.
- Alagić, D., N. Zdolec, B. Njari, I. Filipović, A. Ekert Kabalin, G. Čorić-Alagić, L. Kozačinski (2011): Microbial characterisation of horse meat dry sausage. Veterinarski arhiv. Poslan.
- AOAC (2002): Official methods of Analysis (17th ed.). Washington, DC: Association of Official Analytical Chemists.
- Cocolin, L., R. Urso, K. Rantsiou, G. Comi (2005): Identification, sequencing and characterization of lactic acid bacteria genes responsible for bacteriocin production. Tehnologija mesa 46, 162-172.
- Dobranić, V., B. Njari, A. Večkovec, M. Kadivc (2008): Konjsko meso i Hippophagia. Meso 10, 288-292.
- Dobranić, V., B. Njari, B. Mioković, Ž. Cvrtila Fleck, M. Kadivc (2009): Kemijski sastav konjskog mesa. Meso 11, 32-37.
- Feiner, G. (2006): Raw fermented salami. U: Meat products handbook (Feiner, G., ur.). Woodhead Publishing Limited, Cambridge.
- Hughes, M.C., J.P. Kerry, E.K. Arendt, P.M. Kenneally, P.L.H. McSweeney, E.E. O'Neill (2002): Characterisation of proteolysis during the ripening of semi-dry fermented sausages. Meat Sci. 62, 205-216.
- Kozačinski, L., M. Hadžiosmanović, Ž. Cvrtila Fleck, N. Zdolec, I. Filipović, Z. Kozačinski (2008): Kakvoća trajnih kobasica i češnjovki iz individualnih domaćinstava. Meso 10, 45-52.
- Markov, K., J. Frece, D. Čvek, A. Trontel, A. Slavica, D. Kovačević (2010): Dominantna mikroflora fermentiranih kobasica od konjskog mesa. Meso 12, 217-221.



a,b values marked with the same letter (at same day of maturing) are statistically significantly different ( $p < 0.05$ )

Graph 4 Ammonia content during maturing of horsemeat fermented sausages through three seasons



Graph 5 Ammonia content during maturing of horsemeat fermented sausages through three seasons



Figure 5 Cross-section of horsemeat fermented sausages at the end of maturing

## Qualität der fermentierten Würste aus Pferdefleisch während drei Herstellungssaisons

### Zusammenfassung

Die Herstellung der Würste aus Pferdefleisch hat in unserem Land eine lange Tradition, die unsere gastronomische Tradition bereichert. Es gibt aber zu wenig zugängliche Angaben über diese traditionelle Herstellung und Qualität der Erzeugnisse, um von Standardisation und Schutz des Originalerzeugnisses sprechen zu können. Deshalb war das Ziel dieser Arbeit physikalisch-chemische und sensorische Eigenschaften der Würste aus Pferdefleisch zu untersuchen, dies im Zusammenhang von der Herstellungssaison und Reifephase. Die durchschnittlichen pH-Werte im fertigen Erzeugnis betrugen 5,83, Wasseranteil 19,90 %, Wasseraktivität 0,896, NaCl-Menge 3,44 %, Ammoniakmenge 0,42 mg%. Durch sensorische Bewertung wurden unbedeutende Abweichungen bei sensorischen Eigenschaften der Würste in Bezug auf die Herstellungssaison beobachtet. Während des Reifeprozesses wurden je nach Herstellungssaison statistisch bedeutende Unterschiede ( $p < 0,05$ ) der physikalisch-chemischen Indikatoren festgestellt, die aber trotzdem keinen Einfluss auf die Ausgeglichenheit der Qualität von fertigen Erzeugnissen hatten.

**Schlüsselwörter:** fermentierte Würste, Pferdefleisch, physikalisch-chemische Veränderungen, Qualität

## Qualità delle salsicce fermentate fatte di carne di cavallo nel corso di tre stagioni produttive

### Sommario

La produzione delle salsicce fermentate fatte di carne di cavallo ha una lunga tradizione nel nostro paese il che arricchisce la nostra offerta gastronomica. Ci sono però troppo pochi dati su tale produzione tradizionale e su tale qualità del prodotto per poter parlare della standardizzazione e protezione dell'originalità. Lo scopo di questo lavoro era perciò di fare una ricerca sulle caratteristiche fisico-chimiche e quelle sensoriche delle salsicce fatte di carne di cavallo che dipendevano dalla stagione produttiva e dalla fase di maturazione. I valori pH medi del prodotto finale facevano 5,83, la percentuale dell'acqua faceva 19,90 %, l'attività dell'acqua 0,896, la percentuale del NaCl 3,44 % e dell'ammoniaca 0,42 mg%. La valutazione sensorica ha mostrato le minime differenze nelle caratteristiche sensoriche delle salsicce rispetto alla stagione della produzione. Durante la maturazione le differenze degli indici fisico-chimici notevoli quanto alla statistica ( $p < 0,05$ ) sono state confermate che però non hanno avuto un influsso evidente sull'uguagliamento della qualità del prodotto finale.

**Parole chiave:** salsicce fermentate, carne di cavallo, cambiamenti fisico-chimici, qualità

Njari, B. (2010): Ždrebeća kobasica – eko-proizvod. EKO-VET Fest, Bizovačke Toplice, 18.03.-20.03.2010.

Paleari, M., V. Moretti, G. Beretta, T. Mentasi (2003): Cured products from different animal species. Meat sci. 63, 485-489.

Pleadin, J., N. Peršić, A. Vulić, J. Đugum (2009): Kakvoća trajnih, polutrajnih i obarenih kobasica na hrvatskom tržištu. Hrvatski časopis za prehrambenu tehnologiju, biotehnologiju i nutricionizam 4, 104-108.

Stamenković, T., B. Dević (1990):

Organoleptička svojstva kobasica od konjskog mesa, Tehnologija mesa 2, 77-79.

Šimić, D., B. Mioković (2008): Prilog poznavanju suhih kobasica od konjskog mesa («piketa») iz okolice Pakraca. Meso 10, 292-296.

Zdolec, N., M. Hadžiosmanović, L. Kozačinski, Ž. Cvrtila, I. Filipović, K. Leskovar, N. Vragović, D. Budimir (2007): Fermentirane kobasice proizvedene u domaćinstvu – mikrobiološka kakvoća. Meso 9, 318-324.

Zdolec, N., M. Hadžiosmanović, L.

Kozačinski, Ž. Cvrtila, I. Filipović, M. Škrivanko, K. Leskovar (2008): Microbial and physicochemical succession in fermented sausages produced with bacteriocinogenic culture of *Lactobacillus sakei* and semi-purified bacteriocin mesenterocin Y. Meat Sci. 80, 480-487.

Živković, J. (1986): Higijena i tehnologija mesa II dio. Kakvoća i prerada. "Tipografija" Đakovo.

Received: April 8, 2011

Accepted: June 27, 2011



B | R | N | O

University of Veterinary and Pharmaceutical Sciences Brno  
FACULTY OF VETERINARY HYGIENE AND ECOLOGY  
and State Veterinary Administration of the Czech Republic

## Food Hygiene and Technology

## 41<sup>st</sup> Lenfeld and Hökl's Days

the 250<sup>th</sup> World Anniversary of Veterinary Education

12<sup>th</sup> – 13<sup>th</sup> October 2011

The conference is organized with the financial support of Brno City Municipality.